



Parent/Athlete Concussion Information Sheet

A concussion is a type of traumatic brain injury that changes the way the brain normally works. A concussion is caused by bump, blow, or jolt to the head or body that causes the head and brain to move rapidly back and forth. Even a "ding," "getting your bell rung," or what seems to be a mild bump or blow to the head can be serious.

WHAT ARE THE SIGNS AND SYMPTOMS OF CONCUSSION?

Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury.

If an athlete reports **one or more** symptoms of concussion listed below after a bump, blow, or jolt to

Did You Know?

- Most concussions occur *without* loss of consciousness.
- Athletes who have, at any point in their lives, had a concussion have an increased risk for another concussion.
- Young children and teens are more likely to get a concussion and take longer to recover than adults.

the head or body, s/he should be kept out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it's OK to return to play.

SIGNS OBSERVED BY COACHING STAFF	SYMPTOMS REPORTED BY ATHLETES
Appears dazed or stunned	Headache or "pressure" in head
Is confused about assignment or position	Nausea or vomiting
Forgets an instruction	Balance problems or dizziness
Is unsure of game, score, or opponent	Double or blurry vision
Moves clumsily	Sensitivity to light
Answers questions slowly	Sensitivity to noise
Loses consciousness (<i>even briefly</i>)	Feeling sluggish, hazy, foggy, or groggy
Shows mood, behavior, or personality changes	Concentration or memory problems
Can't recall events <i>prior</i> to hit or fall	Confusion
Can't recall events <i>after</i> hit or fall	Just not "feeling right" or "feeling down"

CONCUSSION DANGER SIGNS

In rare cases, a dangerous blood clot may form on the brain in a person with a concussion and crowd the brain against the skull. An athlete should receive immediate medical attention if after a bump, blow, or jolt to the head or body s/he exhibits any of the following danger signs:

- One pupil larger than the other
- Is drowsy or cannot be awakened
- A headache that not only does not diminish, but gets worse
- Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Cannot recognize people or places
- Becomes increasingly confused, restless, or agitated
- Has unusual behavior
- Loses consciousness (*even a brief loss of consciousness should be taken seriously*)

WHY SHOULD AN ATHLETE REPORT THEIR SYMPTOMS?

If an athlete has a concussion, his/her brain needs time to heal. While an athlete's brain is still healing, s/he is much more likely to have another concussion. Repeat concussions can increase the time it takes to recover. In rare cases, repeat concussions in young athletes can result in brain swelling or permanent damage to their brain. *They can even be fatal.*

It's better to miss one game than the whole season. For more information on concussions, visit: www.cdc.gov/Concussion.

Remember

Concussions affect people differently. While most athletes with a concussion recover quickly and fully, some will have symptoms that last for days, or even weeks. A more serious concussion can last for months or longer.

WHAT SHOULD YOU DO IF YOU THINK YOUR ATHLETE HAS A CONCUSSION?

If you suspect that an athlete has a concussion, remove the athlete from play and seek medical attention. Do not try to judge the severity of the injury yourself. Keep the athlete out of play the day of the injury and until a health care professional, experienced in evaluating for concussion, says s/he is symptom-free and it's OK to return to play.

Rest is key to helping an athlete recover from a concussion. Exercising or activities that involve a lot of concentration, such as studying, working on the computer, or playing video games, may cause concussion symptoms to reappear or get worse. After a concussion, returning to sports and school is a gradual process that should be carefully managed and monitored by a health care professional.

Student-Athlete Name Printed

Student-Athlete Signature

Date

Parent or Legal Guardian Printed

Parent or Legal Guardian Signature

Date

Concussion: Return to Play Progression

Baseline (Step 0): As the baseline step of the Return to Play Progression, the athlete *must have completed total physical and cognitive rest and not be experiencing concussion symptoms for a minimum of 24 hours.* Keep in mind, the younger the athlete, the more conservative the treatment.

Step 1: Light Aerobic Exercise

The Goal: only to increase an athlete's heart rate.

The Time: 5 to 10 minutes.

The Activities: exercise bike, walking, or light jogging.

Absolutely no weight lifting, jumping or hard running.

Step 2: Moderate Exercise

The Goal: limited body and head movement.

The Time: Reduced from typical routine

The Activities: moderate jogging, brief running, moderate-intensity stationary biking, and moderate-intensity weightlifting

Step 3: Non-contact Exercise

The Goal: more intense but non-contact

The Time: Close to Typical Routine

The Activities: running, high-intensity stationary biking, the player's regular weightlifting routine, and non-contact sport-specific drills. This stage may add some cognitive component to practice in addition to the aerobic and movement components introduced in Steps 1 and 2.

Step 4: Practice

The Goal: Reintegrate in full contact practice.

Step 5: Play

The Goal: Return to competition

-
- It is important to monitor symptoms and cognitive function carefully during each increase of exertion.
 - Athletes should only progress to the next level of exertion if they are not experiencing symptoms at the current level. *If symptoms return at any step, an athlete should stop these activities as this may be a sign the athlete is pushing too hard.* Only after additional rest, when the athlete is once again not experiencing symptoms for a minimum of 24 hours, should he or she start again at the previous step during which symptoms were experienced.
 - The Return to Play Progression process is best conducted through a team approach and by a health professional who knows the athlete's physical abilities and endurance. By gauging the athlete's performance on each individual step, a health care professional will be able to determine how far to progress the athlete on a given day. In some cases, the athlete may be able to work through one step in a single day, while in other cases it may take several days to work through an individual step. It may take several weeks to months to work through the entire 5-step progression.